

# THE SEARCH FOR THE SMALLEST 3-E.C. GRAPHS

PRZEMYSŁAW GORDINOWICZ

Theorem

(This is a joint work with Paweł Prałat.)

Adjacency properties of graphs, first studied by Erdős and Rényi, have received much attention. One such property is the  $n$ -e.c. property. For a positive integer  $n$ , a graph is  $n$ -*existentially closed* or  $n$ -*e.c.*, if for all disjoint sets of vertices  $A$  and  $B$  with  $|A \cup B| = n$  (one of  $A$  or  $B$  can be empty), there is a vertex  $z$  not in  $A \cup B$  joined to each vertex of  $A$  and no vertex of  $B$ . We say that  $z$  is *correctly joined* to  $A$  and  $B$ . For completeness, every graph is 0-e.c.

Because for every positive integer  $n$  there is a graph that satisfies  $n$ -e.c. property, therefore we can define  $m_{ec}(n)$  to be the minimum order of an  $n$ -e.c. graph. It is known that  $m_{ec}(1) = 4$  and  $m_{ec}(2) = 9$ , but no other values of this function are determined. For example,  $20 \leq m_{ec}(3) \leq 28$ . Similar to the well-known Ramsey numbers, it is difficult to compute the exact value of  $m_{ec}(n)$ , even for  $n = 3$ , and very little progress has been made.

During my talk, I would like to present a few initial results that improve the lower bound, namely,  $m_{ec}(3) \geq 23$ . An initial results also *suggest* that there is no 3-e.c. graph of order 23 and I hope to be able to announce this during my talk. In order to obtain the result, a computer support was required to check that no graph satisfies a necessary condition described in the following theorem.

**Theorem 1.** *Let  $n \geq m \geq 1$ . If  $G$  is  $n$ -e.c., then for all disjoint sets of vertices  $X$  and  $Y$  with  $|X \cup Y| = m$  (one of  $X$  or  $Y$  can be empty), a graph induced by vertex set  $Z = Z(X, Y)$  defined as*

$$Z = \left( \bigcap_{x \in X} N(x) \right) \cap \left( \bigcap_{y \in Y} N^c(y) \right)$$

*is  $(n - m)$ -e.c.*

The main idea is based on this theorem but an additional interesting and subtle approaches were required to obtain the result.

During my talk, I would like to also describe a few open problems in this field, including the following intriguing one. From the above

theorem we have that  $m_{ec}(n) \geq 2m_{ec}(n-1) + 1$ . As  $m_{ec}(3) \geq 23$  we derive that  $m_{ec}(n) \geq 3 \cdot 2^n - 1$ . On the other hand, using the random graph  $G(v, 1/2)$  one has that  $m_{ec}(n) = O(n^2 2^n)$ . The most important open problem in this area is to determine whether  $\lim_{n \rightarrow \infty} \frac{m_{ec}(n)}{2^n}$  exists and, if so to find its value.

INSTITUTE OF MATHEMATICS, TECHNICAL UNIVERSITY OF ŁÓDŹ, POLAND  
*E-mail address:* `pgordin@p.lodz.pl`